

Hemispheric Differences in Tropical Lower Stratospheric Transport and Tracers Annual Cycle

A11U: Processes and Linkages in the Upper Troposphere and Lower Stratosphere: Observations and Models I

Olga Tweedy¹, D. Waugh¹, R. Stolarski^{1,2}, L. Oman²

1. Johns Hopkins University, Department of Earth and Planetary Sciences;
2. NASA Goddard Space Flight Center

Transport: Vertical advection vs quasi-horizontal mixing

Olga Tweedy

Introduction


Data

Results

Conclusions

The balance between upwelling and quasi-horizontal mixing in the tropical lower stratosphere is not well understood!

- Randel *et al.* [2007]: large seasonal cycle in upwelling is a response for seasonal changes in ozone.
- Abalos *et al.* [2012, 2013] and Ploeger *et al.* [2012] show the importance of eddy mixing



These studies have focused on variations in tracers and processes in the tropic-wide average (20°N-20°S), i.e. have considered “well-mixed” tropics.

Hemispheric differences in ozone annual cycle

Olga Tweedy

Introduction

Data



Results

Conclusions

Southern tropics (ST) are different from Northern tropics (NT)

- Larger annual amplitude in NT
- 2-3 month shift in phase.

[Stolarski et al., 2014]



Annual cycle amplitude of ozone (% relative to the mean) from satellite observations

MAIN QUESTIONS

Olga Tweedy Introduction Data Results Conclusions

1. Do Chemistry Climate Models (CCMs) capture the observed differences between the NT and ST?
2. What processes control the ST and NT annual cycles in the models?

CCMVal -2 MODELS

Olga Tweedy

Introduction

Data

Results

Conclusions

Model Simulations: 1960 to 2010 simulation of the models from CCMVal-2 multi-model intercomparison project

- ❖ 18 CCMs: all transient (historical) runs with nearly identical forcings (GHGs, ODSs, ect).

Satellite observations:

- ❖ Version 3 and 4 of the Aura Microwave Limb Sounder (**MLS**) [Livesey et al., 2008]
- ❖ the Ozone Mapping and Prole Suite (**OMPS**) on board NASA/NOAA Suomi-NPP satellite [Kramarova et al., 2014]
- ❖ the Stratospheric Aerosol and Gas Experiment II (**SAGEII**) [Wang et al., 2002]

Latitude-pressure variations in ozone seasonality

Olga Tweedy


Introduction

Data


Results

Conclusions

Observations



Models



MULTI-MODEL COMPARISON

Olga Tweedy


Introduction

Data

Results

Conclusions

Do Chemistry Climate Models (CCMs) capture the observed differences between NT and ST?



- Distinguished between **NT** (0-20°N) and **ST** (0-20°S)
- In most of CCMVal-2 models NT amplitude > ST amplitude.
- Large spread in amplitudes among models.

Does seasonality in upwelling explain seasonality in ozone?


Olga Tweedy

Introduction

Data and Methods

Results

Conclusions



ST: spread in w^* amplitude among CCMVal-2 models “explains” most of spread in O_3 amplitude

Conclusion: upwelling is controlling factor in the **ST**



Range of amplitudes in O_3 from observations and w^* from reanalysis

Does seasonality in upwelling explain seasonality in ozone?

Olga Tweedy

Introduction


Data and Methods

Results

Conclusions

NT: no relationship between w^* amplitude and O_3 amplitude among CCMVal-2 models.

Conclusion: other factors determine ozone seasonality (mixing may be more important)



Quantifying transport affects on ozone seasonality

Olga Tweedy

Introduction

Data

Results

Conclusions

1. Two models from CCMVal-2 project: **WACCM** and **GEOSCCM**
2. Distinguished between **NT** (0- 18°N) and **ST**(0-18°S)
3. Transform Eulerian Mean analysis (**TEM**) [Andrews, 1987]:

$$\bar{\chi}_t = \left[-\bar{v}^* \bar{\chi}_y + e^{z/H} (\cos \varphi)^{-1} (M_y \cos \varphi)_y \right] + \left[-\bar{w}^* \bar{\chi}_z + e^{z/H} (M_z)_z \right] + P - L.$$

-to isolate role of vertical transport (**red**), horizontal transport (**green**), and chemistry (**blue**).

Processes controlling the NT and ST annual cycles


Olga Tweedy

Introduction

Data

Results

Conclusions



❖ **NT** – horizontal mixing is a dominant cause of seasonality

❖ **ST** – upwelling is a dominant cause of seasonality

$$\bar{\chi}_t = \underbrace{\left[-\bar{v}^* \bar{\chi}_y + e^{z/H} (\cos \varphi)^{-1} (M_y \cos \varphi)_y \right]}_{\text{green line}} + \underbrace{\left[-\bar{w}^* \bar{\chi}_z + e^{z/H} (M_z)_z \right]}_{\text{red line}} + \underbrace{P - L}_{\text{blue line}}.$$

Zonal variations in annual cycle amplitude

Olga Tweedy


Introduction

Data



Results

Conclusions




80hPa July






80hPa January



❖ **July:** Inflow of ozone rich air from NH extratropics into the tropics by N. American and Asian Summer Monsoon



❖ **January:** Ozone is mixed into the tropics over Atlantic and Pacific oceans by Rossby wave breaking



CONCLUSIONS

Olga Tweedy

Introduction

Data

Results

Conclusions

- ❖ Paradigm of well-mixed tropics have to be reconsidered
- ❖ The majority of the CCMs produced the observed feature of a larger annual cycle in the NT than ST
- ❖ The spread among the models much larger than in observations suggesting large differences in transport among the models
- ❖ NT-ST contrast is due to differences in balance between transport processes:
 - ❖ Seasonality in upwelling is most important in the ST
 - ❖ Seasonality in horizontal mixing – in the NT.



ACKNOWLEDGEMENTS



This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program (NSF GRFP)

Authors are grateful to M. Abalos for providing them with the WACCM data

Thank you! Questions?



JOHNS HOPKINS
UNIVERSITY

REFERENCES

- Garcia *et al.* (2007), Simulation of secular trends in the middle atmosphere, J. Geophys. Res., 112, D09301, doi:10.1029/2006JD007485
- Konopka *et al.* (2010), Annual cycle of ozone at and above the tropical tropopause: Observations versus simulations with the Chemical Lagrangian Model of the Stratosphere (CLaMS), Atmos. Chem. Phys., 10(1), 121–132, doi:10.5194/acp-10-121-2010
- Pawson, *et al.* (2008), Goddard Earth Observing System chemistry-climate model simulations of stratospheric ozone-temperature coupling between 1950 and 2005, J. Geophys. Res., 113, D12103, doi:10.1029/2007JD009511.
- Stolarski *et al.* (2014), Seasonal variation of ozone in the tropical lower stratosphere: Southern tropics are different from northern tropics, J. Geophys. Res. Atmos., 119, doi: 10.1002/2013JD021294
- Andrews, D. G., Holton, J. R., and Leovy, C. B.: Middle Atmosphere Dynamics, Academic Press, Orlando, Florida, 489 pp, 1987